

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	for inhibiting and treating bone loss by peptides					Go	Clear
Limits Preview/Index History Clipboard							

Entrez
PubMed

Display	Summary	Save	Text	Order	Details	Add to Clipboard
Show: 20	Items 1-12 of 12					One page.


PubMed
ServicesRelated
Resources

- ☐ **1:** Feige U, Hu YL, Gasser J, Campagnuolo G, Muniyakazi L, Bolon B. Related Articles
Anti-interleukin-1 and anti-tumor necrosis factor-alpha synergistically inhibit adjuvant arthritis in Lewis rats.
Cell Mol Life Sci. 2000 Sep;57(10):1457-70.
PMID: 11078023 [PubMed - indexed for MEDLINE]
- ☐ **2:** Affinito P, Palomba S, Sorrentino C, Pellicano M, Morgera R, Nappi C. Related Articles
[Postmenopausal osteoporosis: therapeutic approaches].
Minerva Ginecol. 1997 Mar;49(3):109-20. Review. Italian.
PMID: 9173338 [PubMed - indexed for MEDLINE]
- ☐ **3:** Funaba M, Murata T, Murata E, Abe M, Takahashi M, Torii K. Related Articles
Increased cartilage and bone formation in spontaneously hypercholesterolemic rats.
Life Sci. 1997;61(6):645-52.
PMID: 9250721 [PubMed - indexed for MEDLINE]
- ☐ **4:** Kimble RB, Matayoshi AB, Vannice JL, Kung VT, Williams C, Pacifici R. Related Articles
Simultaneous block of interleukin-1 and tumor necrosis factor is required to completely prevent bone loss in the early postovariectomy period.
Endocrinology. 1995 Jul;136(7):3054-61.
PMID: 7789332 [PubMed - indexed for MEDLINE]
- ☐ **5:** Prestwood KM, Pilbeam CC, Raisz LG. Related Articles
Treatment of osteoporosis.
Annu Rev Med. 1995;46:249-56. Review.
PMID: 7598461 [PubMed - indexed for MEDLINE]
- ☐ **6:** Polatti F, Perotti F, Angelini GP, Vassellatti D, Rapisardi I. Related Articles
Effects of salmon calcitonin suppositories in the prevention of bone loss in oophorectomized women.
Maturitas. 1993 Dec;18(1):73-6.
PMID: 8107619 [PubMed - indexed for MEDLINE]
- ☐ **7:** Belch AR, Bergsagel DE, Wilson K, O'Reilly S, Wilson J, Sutton D, Pater J, Johnston D, Zee B. Related Articles
Effect of daily etidronate on the osteolysis of multiple myeloma.
J Clin Oncol. 1991 Aug;9(8):1397-402.
PMID: 1712835 [PubMed - indexed for MEDLINE]

- ☐ **8:** [Liu CC, Kalu DN.](#) Related Articles
Human parathyroid hormone-(1-34) prevents bone loss and augments bone formation in sexually mature ovariectomized rats.
J Bone Miner Res. 1990 Sep;5(9):973-82.
PMID: 2281827 [PubMed - indexed for MEDLINE]
- ☐ **9:** [Haas HG, Liebrich BM, Schaffner W.](#) Related Articles
[Calcitonin and osteoporosis--a critical review of the literature 1980-1989].
Klin Wochenschr. 1990 Apr 2;68(7):359-71. Review. German.
PMID: 2188043 [PubMed - indexed for MEDLINE]
- ☐ **10:** [Reginster JY, Deroisy R, Lecart MP, Sarlet N, Fontaine MA, Albert A, Franchimont P.](#) Related Articles
Calcitonin and postmenopausal bone loss.
Exp Gerontol. 1990;25(3-4):349-56.
PMID: 2226670 [PubMed - indexed for MEDLINE]
- ☐ **11:** [Di Silverio F, Serio M, D'Eramo G, Sciarra F.](#) Related Articles
Zoladex vs. Zoladex plus cyproterone acetate in the treatment of advanced prostatic cancer: a multicenter Italian study.
Eur Urol. 1990;18 Suppl 3:54-61.
PMID: 2151278 [PubMed - indexed for MEDLINE]
- ☐ **12:** [MacIntyre I, Stevenson JC, Whitehead MI, Wimalawansa SJ, Banks LM, Healy MJ.](#) Related Articles
Calcitonin for prevention of postmenopausal bone loss.
Lancet. 1988 Apr 23;1(8591):900-2.
PMID: 2895829 [PubMed - indexed for MEDLINE]

Display	Summary	<input checked="" type="checkbox"/>	Save	Text	Order	Details	Add to Clipboard
Show:	20	<input checked="" type="checkbox"/>	Items 1-12 of 12				One page.

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	for TRANCE/RANK INHIBITOR					Go	Clear
Limits Preview/Index History Clipboard							

Entrez
PubMed

Display	Summary	Save	Text	Order	Details	Add to Clipboard
Show: 20	Items 1-4 of 4					One page.

NEW

PubMed
Services

- ☐ 1: [Atkins GJ, Haynes DR, Graves SE, Evdokiou A, Hay S, Bouralexis S, Findlay DM.](#) Related Articles

Expression of osteoclast differentiation signals by stromal elements of giant cell tumors.

J Bone Miner Res. 2000 Apr;15(4):640-9.

PMID: 10780856 [PubMed - indexed for MEDLINE]

- ☐ 2: [Chen NJ, Huang MW, Hsieh SL.](#) Related Articles

Enhanced secretion of IFN-gamma by activated Th1 cells occurs via reverse signaling through TNF-related activation-induced cytokine.

J Immunol. 2001 Jan 1;166(1):270-6.

PMID: 11123302 [PubMed - indexed for MEDLINE]

Related
Resources

- ☐ 3: [Roodman GD.](#) Related Articles

Cell biology of the osteoclast.

Exp Hematol. 1999 Aug;27(8):1229-41. Review.

PMID: 10428500 [PubMed - indexed for MEDLINE]

- ☐ 4: [Wong BR, Josien R, Lee SY, Vologodskaja M, Steinman RM, Choi Y.](#) Related Articles, Protein, Nucleotide

The TRAF family of signal transducers mediates NF-kappaB activation by the TRANCE receptor.

J Biol Chem. 1998 Oct 23;273(43):28355-9.

PMID: 9774460 [PubMed - indexed for MEDLINE]

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

Institution: USPTO || [Sign In as Member or Individual \(Non-member\)](#) || [Contact Subscription Administrator at this Site](#) || [FAQ](#)

The Journal of Immunology, 2001, 166: 270-276.
Copyright © 2001 by [The American Association of Immunologists](#)

Enhanced Secretion of IFN- γ by Activated Th1 Cells Occurs Via Reverse Signaling Through TNF-Related Activation-Induced Cytokine¹

Nien-Jung Chen^{*}, Mei-Wen Huang^{*} and Shie-Liang Hsieh^{2,*†}

^{*} Department of Microbiology and Immunology, and [†] Immunology Research Center, National Yang-Ming University, Taipei, Taiwan


- ▶ [Abstract of this Article](#)
- ▶ [Reprint \(PDF\) Version of this Article](#)
- ▶ Similar articles found in:
 [The JI](#)
 [PubMed](#)
- ▶ [PubMed Citation](#)
- ▶ Search Medline for articles by:
 [Chen, N.-J.](#) || [Hsieh, S.-L.](#)
- ▶ Alert me when:
 [new articles cite this article](#)
- ▶ [Download to Citation Manager](#)

Abstract

Growing evidence has demonstrated that members of TNF superfamily transduce signals after engagement with their receptors. TNF-related activation-induced cytokine (TRANCE), a member of TNF superfamily, is preferentially expressed on the surface of activated CD4⁺ Th1 cells. The soluble receptor activator of NF- κ B (RANK).Fc fusion protein suppresses IFN- γ secretion by activated Th1 cells, but does not affect IL-4 secretion by Th2 cells. The suppressive effect on IFN- γ secretion is observed when Th1 cells are activated by APCs, but not by immobilized anti-TCR β mAb. In contrast, immobilized RANK.Fc fusion protein augments IFN- γ secretion by Th1 cells, indicating the occurrence of reverse signaling through TRANCE during T cell/APC interaction. The enhanced secretion of IFN- γ mediated via TRANCE correlates with the activation of p38 mitogen-activated protein kinase and is blocked by SB203580, a p38 mitogen-activated protein kinase-specific inhibitor. Thus, in addition to its role in activating dendritic cells by binding to the receptor RANK, TRANCE itself can signal the augmentation of IFN- γ secretion via a p38-dependent pathway, and this provides yet another example of reverse signaling by a member of TNF superfamily.

- ▲ [Top](#)
- [Abstract](#)
- ▼ [Introduction](#)
- ▼ [Materials and Methods](#)
- ▼ [Results](#)
- ▼ [Discussion](#)
- ▼ [References](#)

Introduction



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	▼	for		Go	Clear		
Limits		Preview/Index		History		Clipboard	

Entrez
PubMed

Display	Abstract ▼	Save	Text	Order	Add to Clipboard
---------	------------	------	------	-------	------------------

☐ 1: J Bone Miner Res 2000 Apr;15(4):640-9

Related Articles, Books, LinkOut

NEW

Expression of osteoclast differentiation signals by stromal elements of giant cell tumors.

PubMed
Services


Atkins GJ, Haynes DR, Graves SE, Evdokiou A, Hay S, Bouralexis S, Findlay DM.

Department of Orthopaedics and Trauma, University of Adelaide, South Australia.

Related
Resources

The mechanisms by which primary tumors of the bone cause bone destruction have not been elucidated. Unlike most other lytic bone tumors, osteoclastomas, otherwise known as giant cell tumors (GCT), contain osteoclast-like cells within the tumor stroma. A new member of the TNF-ligand superfamily member, osteoclast differentiation factor (ODF/OPGL/RANKL/TRANCE), was recently identified. ODF was shown to directly stimulate osteoclastogenesis, in the presence of M-CSF. In this study, the expression of ODF was examined in a number of tumor samples associated with bone lysis in vivo. In addition, we investigated expression of the ODF receptor on osteoclast precursors, RANK, as well as the ODF inhibitor osteoprotegerin (OPG), and another TNF-ligand superfamily member, TRAIL, previously shown to abrogate the inhibitory effects of OPG. We report here the novel finding that GCT stromal cells contain abundant ODF mRNA, whereas the giant cell population exclusively expresses RANK mRNA. These results are consistent with the osteoclast-mediated bone destruction by these tumors. We also report the expression of OPG and TRAIL mRNA in GCT samples. A comparison with other lytic and nonlytic tumors of bone showed that GCT express more ODF and TRAIL mRNA relative to OPG mRNA. In addition, GCT were found to express a number of cytokines previously reported to play central roles in osteoclastogenesis, namely, IL-1, -6, -11, -17, as well as TNF-alpha. Importantly, GCT were also found to express high levels of M-CSF mRNA, a cytokine shown to be an essential cofactor of ODF, and a survival factor for mature and developing osteoclasts. Furthermore, expression of these molecules by stromal cells isolated from GCT continued in vitro. Thus GCT constitutively express all of the signals that are currently understood to be necessary for the differentiation of osteoclasts from precursor cells.

PMID: 10780856 [PubMed - indexed for MEDLINE]



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	
Search	PubMed	▼	for				Go	Clear
Limits		Preview/Index		History		Clipboard		

Entrez
PubMed

☐ 1: Endocrinology 1995 Jul;136(7):3054-61

Related Articles, Books

NEW

PubMed
Services

Simultaneous block of interleukin-1 and tumor necrosis factor is required to completely prevent bone loss in the early postovariectomy period.


Kimble RB, Matayoshi AB, Vannice JL, Kung VT, Williams C, Pacifici R.

Division of Bone and Mineral Diseases, Washington University School of Medicine, St. Louis, Missouri, USA.

Related
Resources

Considerable evidence supports the hypothesis that estrogen prevents bone loss by blocking the production of cytokines in bone or bone marrow. However, controversy remains on the role of candidate factors, such as interleukin-1 (IL-1), IL-6, and tumor necrosis factor (TNF). As IL-1 and TNF have many additive and/or synergistic effects in bone, we tested the hypothesis that the simultaneous block of IL-1 and TNF is required to prevent the initial phase of rapid bone loss that follows ovariectomy (ovx). To this aim, rats were ovariectomized and treated for 2 weeks with either IL-1 receptor antagonist (IL-1ra), an inhibitor of IL-1, or TNF-binding protein (TNFbp), an inhibitor of TNF. Ovx increased bone marrow cell secretion of IL-1 and TNF and decreased the bone density of the distal femur, as measured by dual energy x-ray absorptiometry. Ovx-induced bone loss was decreased by both IL-1ra and TNFbp and completely prevented by simultaneous treatment with IL-1ra and TNFbp. Combined treatment with IL-1ra and TNFbp decreased urinary pyridinoline cross-links, a marker of bone resorption that reflects osteoclast number and osteoclast activity, whereas treatment with either inhibitor alone was less effective. Both IL-1ra and TNFbp decreased the number of osteoclasts on the endocortical surfaces and stimulated bone formation, but the two inhibitors had no additive effects on these indexes, suggesting that inhibition of osteoclastogenesis and stimulation of bone formation do not account for the additive bone-sparing effects of IL-1ra and TNFbp. These inhibitors had no effect in sham-operated rats, indicating that they specifically blocked estrogen-dependent events. In conclusion, these data indicate that in the early post-ovx period, IL-1 and TNF play a critical causal role in inducing bone loss and do so by stimulating bone resorption and inhibiting bone formation.

PMID: 7789332 [PubMed - indexed for MEDLINE]



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	for					Go	Clear
Limits Preview/Index History Clipboard							

Entrez
PubMed

☐ 1: J Clin Periodontol 2001 Mar;28(3):233-40

Related Articles, Books, LinkOut

Online

NEW

Soluble antagonists to interleukin-1 (IL-1) and tumor necrosis factor (TNF) inhibits loss of tissue attachment in experimental periodontitis.

PubMed
Services

Delima AJ, Oates T, Assuma R, Schwartz Z, Cochran D, Amar S, Graves DT.

Department of Periodontology and Oral Biology, Boston University School of Dental Medicine, MA, USA.

Related
Resources

BACKGROUND, AIMS: Periodontal disease is a significant cause of tooth loss among adults and is characterized by the alteration and permanent destruction of the deeper periodontal tissues. Although the presence of pathologic microbes is required to trigger this process, the amplification and progression of the diseased state is believed to rely heavily on the production of host mediators in response to bacteria or their metabolic products. The inflammatory response is effective in preventing large-scale colonization of the gingival tissues by bacteria that lie in close proximity to the tooth surface or within the gingival sulcus. It has been postulated that the host-response in some individuals may lead to an over-reaction to invading oral pathogens resulting in the destruction of periodontal tissues. **METHODS:** Several host-derived mediators are believed to contribute to this response. Two agents considered to be essential in periodontal destruction are interleukin-1 (IL-1) and tumor necrosis factor (TNF). We investigated the role of IL-1 and TNF in the loss of connective tissue attachment in a *Macaca fascicularis* primate model of experimental periodontitis. Silk ligatures impregnated with the periodontal pathogen, *Porphyromonas gingivalis* were wrapped around the posterior teeth and the activity of IL-1 and TNF were inhibited by soluble receptors to these proinflammatory cytokines via local injection into interdental papillae. **RESULTS:** Histomorphometric analysis indicates that IL-1 and TNF antagonists significantly reduced the loss of connective tissue attachment by approximately 51% and the loss of alveolar bone height by almost 91%, both of which were statistically significant. **CONCLUSION:** This investigation demonstrates that the loss of connective tissue attachment and progression of periodontal disease can be retarded by antagonists to specific host mediators such as IL-1 and TNF and may provide a potential treatment modality to combat the disease process.

Soluble antagonists to interleukin-1 (IL-1) and tumor necrosis factor (TNF) inhibits loss of tissue attachment in experimental periodontitis

A. J. Delima¹, T. Oates², R. Assuma¹, Z. Schwartz^{2,3}, D. Cochran², S. Amar¹ and D. T. Graves¹

Abstract

Background, aims:

Periodontal disease is a significant cause of tooth loss among adults and is characterized by the alteration and permanent destruction of the deeper periodontal tissues. Although the presence of pathologic microbes is required to trigger this process, the amplification and progression of the diseased state is believed to rely heavily on the production of host mediators in response to bacteria or their metabolic products. The inflammatory response is effective in preventing large-scale colonization of the gingival tissues by bacteria that lie in close proximity to the tooth surface or within the gingival sulcus. It has been postulated that the host-response in some individuals may lead to an over-reaction to invading oral pathogens resulting in the destruction of periodontal tissues.

Methods:

Several host-derived mediators are believed to contribute to this response. Two agents considered to be essential in periodontal destruction are interleukin-1 (IL-1) and tumor necrosis factor (TNF). We investigated the role of IL-1 and TNF in the loss of connective tissue attachment in a *Macaca fascicularis* primate model of experimental periodontitis. Silk ligatures

Synergy Homepage	►
Your Homepage	►
List of Journals	►
List of Issues	►
Table of Contents	►
View Article (HTML)	►
View Article (PDF)	►
Notify a Colleague	►
Search	►
Add to Favorites	►
Help	►

PubMed Search

Delima, A
Oates, T
Assuma, R
Schwartz, Z
Cochran, D
Amar, S
Graves, D

Keywords

- ☐ periodontal disease
- ☐ inflammatory mediators
- ☐ cytokine
- ☐ bone
- ☐ inflammation
- ☐ osteoclast
- ☐ cytokine antagonist
- ☐ soluble receptors

Database Search

Affiliations

¹ Department of Periodontology and Oral Biology, Boston University School of Dental Medicine; ² Department of Periodontology, University of Texas Health Science Center, San Antonio, USA; ³ Department of Periodontology, Hebrew University Hadassa, Jerusalem, Israel

Correspondents

Address:
Dana Graves
Department of Periodontology and Oral Biology, Room W-202
Boston University Medical Center
700 Albany Street
Boston, MA 02118, USA
Fax: (617) 638 4924
e-mail: dgraves@acs.bu.edu

impregnated with the periodontal pathogen, *Porphyromonas gingivalis* were wrapped around the posterior teeth and the activity of IL-1 and TNF were inhibited by soluble receptors to these proinflammatory cytokines via local injection into interdental papillae.

Results:

Histomorphometric analysis indicates that IL-1 and TNF antagonists significantly reduced the loss of connective tissue attachment by approximately 51% and the loss of alveolar bone height by almost 91%, both of which were statistically significant.

Conclusion:

This investigation demonstrates that the loss of connective tissue attachment and progression of periodontal disease can be retarded by antagonists to specific host mediators such as IL-1 and TNF and may provide a potential treatment modality to combat the disease process.

To cite this article:

Delima, A. J., Oates, T., Assuma, R., Schwartz, Z., Cochran, D., Amar, S. & Graves, D. T.

Soluble antagonists to interleukin-1 (IL-1) and tumor necrosis factor (TNF) inhibits loss of tissue attachment in experimental periodontitis.

Journal Of Clinical Periodontology **28** (3), 233-240.

Available from:

<http://dx.doi.org/10.1034/j.1600-051x.2001.028003233.x>

Full Text (HTML)

Reprint (PDF)



© Blackwell Science and Munksgaard. All rights reserved. Some contents contained herein may be owned by third parties and use is prohibited without express written permission.